

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat Antiserum anti-Mouse IgG (Fc)-unconj., MinX Bo,Ca,Ck,Dg,Ho,Hu,Mo,Rb,Sh,Sw NMB-GAM/IGG(FC)

Article Name	Goat Antiserum anti-Mouse IgG (Fc)-unconj., MinX Bo,Ca,Ck,Dg,Ho,Hu,Mo,Rb,Sh,Sw
Biozol Catalog Number	NMB-GAM/IGG(FC)
Supplier Catalog Number	GAM/IgG(Fc)
Alternative Catalog Number	NMB-GAM/IGG(FC)
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Mouse
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgG (Fc)
Cross-Adsorption (MinX)	Bovine,Feline,Gallus,Canine,Equine,Human,Monkey,Rabbit,Sheep,Porcine
Product Description	The reactivity of the antiserum is directed to the Fc subunit of the IgG molecule. The antiserum contains antibodies to subclass-specific determinants as well as to determinants shared by two or more subclasses of IgG. In immunoelectrophoresis and do...
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. No foreign proteins a
Source	Purified normal IgG, including all known subclasses, isolated from pooled mouse serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. No foreign proteins a
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and