

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat Antiserum anti-Mouse IgG+IgM+IgA (Fc)-unconj., MinX none NMB-GAM/IG(FC)

Article Name	Goat Antiserum anti-Mouse IgG+IgM+IgA (Fc)-unconj., MinX none
Biozol Catalog Number	NMB-GAM/IG(FC)
Supplier Catalog Number	GAM/Ig(Fc)
Alternative Catalog Number	NMB-GAM/IG(FC)
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Mouse
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgG+IgM+IgA (Fc)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fc subunits of the major immunoglobulins. It does not react with any non-Ig protein in mouse serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In precipitating techniques a...
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum.
Source	Purified polyclonal IgG and homogenous IgA and IgM isolated from mouse serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum.
Antibody Type	Secondary Antibody
Application Notes	Immunoprecipitation. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells.