

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat anti Mouse IgG1 IgG2a IgG2b IgG3 IgA IgM (Fc specific), conjugated with Biotin, Clone: [Polyclonal], Monoclonal NMB-GAM/IG(FC)/BIO

Article Name	Goat anti Mouse IgG1 IgG2a IgG2b IgG3 IgA IgM (Fc specific), conjugated with Biotin, Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GAM/IG(FC)/BIO
Supplier Catalog Number	GAM/Ig(Fc)/Bio
Alternative Catalog Number	NMB-GAM/IG(FC)/BIO
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Mouse
Conjugation	Biotin
Format	IgG
Target Specificity	IgG+IgM+IgA (Fc)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fc subunit of the major immunoglobulin isotypes which expresses strict isotypic (class) specificity. In immunoelectrophoresis and double radial immunodiffusion using various antiserum concentrations ...
Clonality	Monoclonal
Clone Designation	[Polyclonal]

Buffer	Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. IgG protein concentration 10 mg/ml. Biotin/IgG protein molar ratio
Source	Purified normal IgG and homogenous IgA and IgM isolated from mouse serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Form	Biotin
Formula	Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. IgG protein concentration 10 mg/ml. Biotin/IgG protein molar ratio
Antibody Type	Secondary Antibody
Application Notes	ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting.