

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat Antiserum anti-Mouse IgG2ab-unconj., MinX none NMB-GAM/IGG2AB

| | |
|----------------------------|--|
| Article Name | Goat Antiserum anti-Mouse IgG2ab-unconj., MinX none |
| Biozol Catalog Number | NMB-GAM/IGG2AB |
| Supplier Catalog Number | GAM/IgG2ab |
| Alternative Catalog Number | NMB-GAM/IGG2AB |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Species Reactivity | Mouse |
| Conjugation | Unconjugated |
| Format | Antiserum |
| Target Specificity | IgG2ab |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | The reactivity of the antiserum is directed to the subclasses IgG2a and IgG2b. The antiserum contains antibodies to iso and allospecific determinants. It does not react with other subclasses of IgG, IgG/Fab fragments, IgM and IgA or any non-Ig protein... |
| Clonality | Polyclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|--|
| Buffer | Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign protei |
| Source | Pools of purified homogenous IgG2a and IgG2b isolated from mouse serum of strains belonging to the allelic types Igh-1a and Igh-1b. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formula | Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign protei |
| Antibody Type | Secondary Antibody |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and |