

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat IgG anti-Mouse IgG (F(ab')<sub>2</sub>)-FITC, MinX Bo,Ca,Ck,Dg,Hm,Ho,Hu,Mo,Rb,Sh,Sw NMB-GAM/FAB/FITC

|                            |   |
|----------------------------|---|
| Article Name               | Goat IgG anti-Mouse IgG (F(ab') <sub>2</sub> )-FITC, MinX<br>Bo,Ca,Ck,Dg,Hm,Ho,Hu,Mo,Rb,Sh,Sw   |
| Biozol Catalog Number      | NMB-GAM/FAB/FITC  |
| Supplier Catalog Number    | GAM/Fab/FITC  |
| Alternative Catalog Number | NMB-GAM/FAB/FITC  |
| Manufacturer               | NordicMubio   |
| Host                       | Goat  |
| Category                   | Antikörper  |
| Application                | ELISA, ICC, IF, IHC-Fr  |
| Species Reactivity         | Mouse   |
| Conjugation                | FITC  |
| Format                     | IgG   |
| Target Specificity         | IgG (F(ab') <sub>2</sub> )  |
| Cross-Adsorption (MinX)    | Bovine,Feline,Gallus,Canine,Hamster<br>(all),Equine,Human,Monkey,Rabbit,Sheep,Porcine   |
| Product Description        | The reactivity of the antiserum is directed to the Fab subunits of intact IgG, IgA IgM and other Ig classes of both light chain types, with their Fab or F(ab) <sub>2</sub> subunits, and with free light chains of kappa and lambda type. In immunoelectrophoresis an... |
| Clonality                  | Polyclonal  |

|                   |  |
|-------------------|--|
| Clone Designation | [Polyclonal]   |
| Buffer            | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun down |
| Source            | Purified Fab of normal IgG isolated from mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.  |
| Form              | FITC   |
| Formula           | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added as it may interfere with the antibody activity.   |
| Antibody Type     | Secondary Antibody   |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.  |