

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat IgG anti-Mouse IgM (μ)-FITC, MinX Hu NMB-GAM/IGM(FC)/FITC

Article Name	Goat IgG anti-Mouse IgM (μ)-FITC, MinX Hu
Biozol Catalog Number	NMB-GAM/IGM(FC)/FITC
Supplier Catalog Number	GAM/IgM(Fc)/FITC
Alternative Catalog Number	NMB-GAM/IGM(FC)/FITC
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Mouse
Conjugation	FITC
Format	IgG
Target Specificity	IgM (μ)
Cross-Adsorption (MinX)	Human
Product Description	The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule. In immunoelectrophoresis and double radial immunodiffusion, using various antiserum concentrations against normal mouse plasma and serum, a single characteristic precipitin arc is observed.
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun down
Source	Purified homogenous IgM isolated from pooled mouse serum. Immunization with intact (19S) and split (7S) IgM. Freunds complete adjuvant is used in the first step of the immunization procedure.
Form	FITC
Formula	FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Antibody Type	Secondary Antibody
Application Notes	ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.