

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Rabbit IgA (Fc specific), Clone: [Polyclonal], Monoclonal NMB-GAR/IGA(FC)

| | |
|----------------------------|--|
| Article Name | Goat anti Rabbit IgA (Fc specific), Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GAR/IGA(FC) |
| Supplier Catalog Number | GAR/IgA(Fc) |
| Alternative Catalog Number | NMB-GAR/IGA(FC) |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Product Description | The reactivity of the antiserum is restricted to the Fc part of the IgA molecule. In immunolectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal rabbit serum a single precipitin line is obtained which shows... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. It is reconstituted by adding 1 |
| Source | Highly purified normal IgA isolated from rabbit serum. Freunds complete adjuvant is used in the first step of the immunization procedure. |

| | |
|-------------------|---|
| Formula | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells. |