

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat Antiserum anti-Rabbit IgG (F(ab')2)-unconj., MinX none NMB-GAR/FAB

| | |
|----------------------------|---|
| Article Name | Goat Antiserum anti-Rabbit IgG (F(ab')2)-unconj., MinX none |
| Biozol Catalog Number | NMB-GAR/FAB |
| Supplier Catalog Number | GAR/Fab |
| Alternative Catalog Number | NMB-GAR/FAB |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Species Reactivity | Rabbit |
| Conjugation | Unconjugated |
| Format | Antiserum |
| Target Specificity | IgG (F(ab')2) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | The antiserum is reacting with the Fab subunit of intact IgG, IgA and IgM and other Ig classes of both light chain types, with their Fab or F(ab)2 subunits and with free light chains of kappa and lambda type as tested in immunoelectrophoresis and dou... |
| Clonality | Polyclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|---|
| Buffer | Delipidated, heat inactivated lyophilized stable whole serum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. It is reconstituted by adding 1 ml s |
| Source | Purified Fab of normal rabbit IgG. Freunds complete adjuvant is used in the first step of the immunization procedure. |
| Formula | Delipidated, heat inactivated lyophilized stable whole serum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. |
| Antibody Type | Secondary Antibody |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 |