

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Rabbit IgG (Fc)-unconj., MinX none NMB-GAR/IGG(FC)/7S

| | |
|----------------------------|---|
| Article Name | Goat IgG anti-Rabbit IgG (Fc)-unconj., MinX none |
| Biozol Catalog Number | NMB-GAR/IGG(FC)/7S |
| Supplier Catalog Number | GAR/IgG(Fc)/7S |
| Alternative Catalog Number | NMB-GAR/IGG(FC)/7S |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Species Reactivity | Rabbit |
| Conjugation | Unconjugated |
| Format | IgG |
| Target Specificity | IgG (Fc) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | The reactivity of the antiserum is directed to the Fc subunit of the IgG molecule which expresses The reactivity of the antiserum is directed to the Fc subunit of the IgG molecule which expresses tested by immunoelectrophoresis and double radial immu... |
| Clonality | Polyclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|--|
| Buffer | Purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove i |
| Source | Purified normal IgG isolated from pooled rabbit serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formula | Purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Antibody Type | Secondary Antibody |
| Application Notes | Indirect immunofluorescence,ELISA,Dot blot,Immunoblotting. |