

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Rabbit IgM (μ)-HRPO, MinX none NMB-GAR/IGM(FC)/PO

| | |
|----------------------------|---|
| Article Name | Goat IgG anti-Rabbit IgM (μ)-HRPO, MinX none |
| Biozol Catalog Number | NMB-GAR/IGM(FC)/PO |
| Supplier Catalog Number | GAR/IgM(Fc)/PO |
| Alternative Catalog Number | NMB-GAR/IGM(FC)/PO |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Species Reactivity | Rabbit |
| Conjugation | HRPO |
| Format | IgG |
| Target Specificity | IgM (μ) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in rabbit serum, as tested by immunoelectrophoresis and double radial imm... |
| Clonality | Polyclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|--|
| Buffer | Horseradish peroxidase-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distill |
| Source | Purified IgM isolated from rabbit serum. Freunds complete adjuvant is used in the first step of the immunization procedure. |
| Form | HRP |
| Antibody Type | Secondary Antibody |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting. |