

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Hamster (all) IgG (H+L)-HRPO, MinX none NMB-GAHA/IGG(H+L)/PO

Article Name	Goat IgG anti-Hamster (all) IgG (H+L)-HRPO, MinX none
Biozol Catalog Number	NMB-GAHA/IGG(H+L)/PO
Supplier Catalog Number	GAHa/IgG(H+L)/PO
Alternative Catalog Number	NMB-GAHA/IGG(H+L)/PO
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Application	DOT, ELISA, ICC, WB
Species Reactivity	Hamster (all)
Conjugation	HRPO
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in hamster serum,
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	Peroxidase-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, sp
Source	Goats were immunized with purified normal IgG isolated from pooled hamster serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	Peroxidase-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Antibody Type	Secondary Antibody
Application Notes	ELISA,Immunocytochemistry,Immunohistochemistry (paraffin),Dot blot,Immunoblotting.