

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat Antiserum anti-Mouse IgG+IgM+IgA+IgD (H+L)-unconj., MinX none NMB-GAM/IG(GAMD)

| | |
|----------------------------|---|
| Article Name | Goat Antiserum anti-Mouse IgG+IgM+IgA+IgD (H+L)-unconj., MinX none |
| Biozol Catalog Number | NMB-GAM/IG(GAMD) |
| Supplier Catalog Number | GAM/Ig(GAMD) |
| Alternative Catalog Number | NMB-GAM/IG(GAMD) |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Species Reactivity | Mouse |
| Conjugation | Unconjugated |
| Format | Antiserum |
| Target Specificity | IgG+IgM+IgA+IgD (H+L) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | The reactivity of the antiserum is directed to the heavy chains of immunoglobulin isotypes IgG, IgA, IgM and IgD. The antiserum also contains antibodies to common determinants and to the surface determinants of the common Fab as tested by immunoelect... |
| Clonality | Polyclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|--|
| Buffer | Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. No foreign proteins a |
| Source | Purified polyclonal mouse IgG and pools of homogenous IgG, IgA, IgM and IgD. Freunds complete adjuvant is used in the first step of the immunization procedure. |
| Formula | Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. No foreign proteins a |
| Antibody Type | Secondary Antibody |
| Application Notes | Immunoprecipitation. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 m |