

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit Antiserum anti-Rat IgG (Fc)-unconj., MinX none NMB-RARA/IGG(FC)

|                            |   |
|----------------------------|---|
| Article Name               | Rabbit Antiserum anti-Rat IgG (Fc)-unconj., MinX none   |
| Biozol Catalog Number      | NMB-RARA/IGG(FC)  |
| Supplier Catalog Number    | RARA/IgG(Fc)  |
| Alternative Catalog Number | NMB-RARA/IGG(FC)  |
| Manufacturer               | NordicMubio   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Species Reactivity         | Rat   |
| Conjugation                | Unconjugated  |
| Format                     | Antiserum   |
| Target Specificity         | IgG (Fc)  |
| Cross-Adsorption (MinX)    | no cross-adsorbtion   |
| Product Description        | The reactivity of the antiserum is directed to the Fc subunit of the IgG molecule. The antiserum contains antibodies to subclass-specific determinants as well as to determinants shared by two or more subclasses of IgG. In immunoelectrophoresis and do... |
| Clonality                  | Polyclonal  |
| Clone Designation          | [Polyclonal]  |

|                   |   |
|-------------------|---|
| Buffer            | Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit s                    |
| Source            | Purified normal IgG, including all known subclasses, isolated from pooled rat serum. Freunds complete adjuvant is used in the first step of the immunization procedure.   |
| Formula           | Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit s                    |
| Antibody Type     | Secondary Antibody  |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l antiserum in a 3 mm diameter centre well and 2 $\mu$ l serum samples (neat and |