

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit Antiserum anti-Rat IgG (H+L)-unconj., MinX Ca,Ck,Sh NMB-RARA/IGG(H+L)

|                            |   |
|----------------------------|---|
| Article Name               | Rabbit Antiserum anti-Rat IgG (H+L)-unconj., MinX Ca,Ck,Sh  |
| Biozol Catalog Number      | NMB-RARA/IGG(H+L)   |
| Supplier Catalog Number    | RARa/IgG(H+L)   |
| Alternative Catalog Number | NMB-RARA/IGG(H+L)   |
| Manufacturer               | NordicMubio   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Species Reactivity         | Rat   |
| Conjugation                | Unconjugated  |
| Format                     | Antiserum   |
| Target Specificity         | IgG (H+L)   |
| Cross-Adsorption (MinX)    | Feline,Gallus,Sheep   |
| Product Description        | The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in rat serum, as t... |
| Clonality                  | Polyclonal  |
| Clone Designation          | [Polyclonal]  |

|                   |  |
|-------------------|--|
| Buffer            | Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit s |
| Source            | Purified normal IgG, including all known subclasses, isolated from pooled rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.   |
| Formula           | Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit s |
| Antibody Type     | Secondary Antibody   |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2  |