

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit IgG anti-Rat IgG (H+L)-FITC, MinX Ck NMB-RARA/IGG(H+L)/FITC

| | |
|----------------------------|---|
| Article Name | Rabbit IgG anti-Rat IgG (H+L)-FITC, MinX Ck |
| Biozol Catalog Number | NMB-RARA/IGG(H+L)/FITC |
| Supplier Catalog Number | RARa/IgG(H+L)/FITC |
| Alternative Catalog Number | NMB-RARA/IGG(H+L)/FITC |
| Manufacturer | NordicMubio |
| Host | Rabbit |
| Category | Antikörper |
| Species Reactivity | Rat |
| Conjugation | FITC |
| Format | IgG |
| Target Specificity | IgG (H+L) |
| Cross-Adsorption (MinX) | Gallus |
| Product Description | The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in rat serum, as t... |
| Clonality | Polyclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|--|
| Buffer | FITC-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added IgG protein concentration 10 mg/ml. Fluorochrome/IgG protein molar ratio (F/P) is approximately 1.3. No foreign pro |
| Source | Purified normal IgG isolated from pooled rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Form | FITC |
| Formula | FITC-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added IgG protein concentration 10 mg/ml. Fluorochrome/IgG protein molar ratio (F/P) is approximately 1.3. No foreign pro |
| Antibody Type | Secondary Antibody |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence. |