

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Bax Antibody, Rabbit, Polyclonal NSJ-F40098-0.08ML

Article Name Bax Antibody, Rabbit, Polyclonal Biozol Catalog Number NSJ-F40098-0.08ML Supplier Catalog Number F40098-0.08ML Alternative Catalog Number NSJ-F40098-0.08ML Manufacturer NSJ Bioreagents Host Rabbit Category Antikörper Application ELISA, FACS, IHC, WB Species Reactivity Human Immunogen Aportion of amino acids 41-76 (BH3 domain) from the human protein was used as the immunogen for this Bax antibody. Product Description bax belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Bax forms a heterodimer with BCL2, and functions as an apo Clonality Polyclonal Concentration In 1X PBS pH 7.4 with 0.09% sodium azide Buffer In 1X PBS pH 7.4 with 0.09% sodium azide		
Supplier Catalog Number F40098-0.08ML Alternative Catalog Number NSJ-F40098-0.08ML Manufacturer NSJ Bioreagents Host Rabbit Category Antikörper Application ELISA, FACS, IHC, WB Species Reactivity Human Immunogen A portion of amino acids 41-76 (BH3 domain) from the human protein was used as the immunogen for this Bax antibody. Product Description Bax belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Bax forms a heterodimer with BCL2, and functions as an apo Clonality Polyclonal Concentration In 1X PBS pH 7.4 with 0.09% sodium azide Isotype Rabbit Ig UniProt Q07812	Article Name	Bax Antibody, Rabbit, Polyclonal
Alternative Catalog Number NSJ-F40098-0.08ML Manufacturer NSJ Bioreagents Host Rabbit Category Antikörper Application ELISA, FACS, IHC, WB Species Reactivity Human Immunogen A portion of amino acids 41-76 (BH3 domain) from the human protein was used as the immunogen for this Bax antibody. Product Description Bax belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Bax forms a heterodimer with BCL2, and functions as an apo Clonality Polyclonal Concentration In 1X PBS pH 7.4 with 0.09% sodium azide Isotype Rabbit Ig UniProt O07812	Biozol Catalog Number	NSJ-F40098-0.08ML
Manufacturer NSJ Bioreagents Host Rabbit Category Antikörper Application ELISA, FACS, IHC, WB Species Reactivity Human A portion of amino acids 41-76 (BH3 domain) from the human protein was used as the immunogen for this Bax antibody. Bax belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Bax forms a heterodimer with BCL2, and functions as an apo Clonality Polyclonal Concentration In 1X PBS pH 7.4 with 0.09% sodium azide Isotype Rabbit Ig UniProt Q07812	Supplier Catalog Number	F40098-0.08ML
Host Rabbit Category Antikörper Application ELISA, FACS, IHC, WB Species Reactivity Human A portion of amino acids 41-76 (BH3 domain) from the human protein was used as the immunogen for this Bax antibody. Bax belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Bax forms a heterodimer with BCL2, and functions as an apo Clonality Polyclonal Concentration In 1X PBS pH 7.4 with 0.09% sodium azide Isotype Rabbit Ig UniProt Q07812	Alternative Catalog Number	NSJ-F40098-0.08ML
Category Antikörper Application ELISA, FACS, IHC, WB Species Reactivity Human Immunogen A portion of amino acids 41-76 (BH3 domain) from the human protein was used as the immunogen for this Bax antibody. Bax belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Bax forms a heterodimer with BCL2, and functions as an apo Clonality Polyclonal Concentration In 1X PBS pH 7.4 with 0.09% sodium azide Isotype Rabbit Ig UniProt Q07812	Manufacturer	NSJ Bioreagents
Application ELISA, FACS, IHC, WB Species Reactivity Human Immunogen A portion of amino acids 41-76 (BH3 domain) from the human protein was used as the immunogen for this Bax antibody. Bax belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Bax forms a heterodimer with BCL2, and functions as an apo Clonality Polyclonal Concentration In 1X PBS pH 7.4 with 0.09% sodium azide Isotype Rabbit Ig UniProt Q07812	Host	Rabbit
Species Reactivity Human A portion of amino acids 41-76 (BH3 domain) from the human protein was used as the immunogen for this Bax antibody. Bax belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Bax forms a heterodimer with BCL2, and functions as an apo Clonality Polyclonal Concentration In 1X PBS pH 7.4 with 0.09% sodium azide UniProt Q07812	Category	Antikörper
Immunogen A portion of amino acids 41-76 (BH3 domain) from the human protein was used as the immunogen for this Bax antibody. Bax belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Bax forms a heterodimer with BCL2, and functions as an apo Clonality Polyclonal Concentration In 1X PBS pH 7.4 with 0.09% sodium azide Isotype Rabbit Ig UniProt Q07812	Application	ELISA, FACS, IHC, WB
Product Description Bax belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Bax forms a heterodimer with BCL2, and functions as an apo Clonality Polyclonal Concentration In 1X PBS pH 7.4 with 0.09% sodium azide Isotype Rabbit Ig UniProt Q07812	Species Reactivity	Human
Product Description hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Bax forms a heterodimer with BCL2, and functions as an apo Clonality Polyclonal Concentration In 1X PBS pH 7.4 with 0.09% sodium azide Isotype Rabbit Ig UniProt Q07812	Immunogen	·
Concentration In 1X PBS pH 7.4 with 0.09% sodium azide Isotype Rabbit Ig UniProt Q07812	Product Description	hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Bax forms a
Isotype Rabbit Ig UniProt Q07812	Clonality	Polyclonal
UniProt Q07812	Concentration	In 1X PBS pH 7.4 with 0.09% sodium azide
	Isotype	Rabbit Ig
Buffer In 1X PBS pH 7.4 with 0.09% sodium azide	UniProt	Q07812
	Buffer	In 1X PBS pH 7.4 with 0.09% sodium azide

Purity	Purified
Form	In 1X PBS pH 7.4 with 0.09% sodium azide
Target	Bax
Antibody Type	Primary Antibody
Application Dilute	Western blot: 1:1000,IHC (Paraffin): 1:50-1:100,Flow Cytometry: 1:10-1:50
Application Notes	Titration of the Bax antibody may be required due to differences in protocols and secondary/substrate sensitivity.