

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Glypican-3 Antibody, Clone: [SGPN3-1], Mouse, Monoclonal NSJ-V7027SAF-100UG

Article Name	Glypican-3 Antibody, Clone: [SGPN3-1], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V7027SAF-100UG
Supplier Catalog Number	V7027SAF-100UG
Alternative Catalog Number	NSJ-V7027SAF-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	FACS, IF, IHC-P
Species Reactivity	Human, Rat
Immunogen	Recombinant full-length human GPC3 protein was used as the immunogen for the Glypican-3 antibody.
Product Description	Glypican-3 is a proteoglycan that bears heparan sulfate. Inhibits the dipeptidyl peptidase activity of DPP4. May be involved in the suppression/modulation of growth in the predominantly mesodermal tissues and organs. May play a role in the modulation...
Clonality	Monoclonal
Concentration	1 mg/ml in 1X PBS, BSA free, sodium azide free
Clone Designation	[SGPN3-1]
Isotype	Mouse IgG1, kappa
UniProt	P51654

Buffer	1 mg/ml in 1X PBS, BSA free, sodium azide free
Purity	Protein G affinity chromatography
Form	1 mg/ml in 1X PBS, BSA free, sodium azide free
Target	Glypican 3
Antibody Type	Primary Antibody
Application Dilute	Flow cytometry: 1-2ug/million cells,Immunofluorescence: 1-2ug/ml,Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT (1),Prediluted IHC only format: incubate for 30 min at RT (2)
Application Notes	Optimal dilution of the Glypican-3 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.