

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin 8 Antibody, Clone: [CYKN8-1], Mouse, Monoclonal NSJ-V7042SAF-100UG

| Article Name | Cytokeratin 8 Antibody, Clone: [CYKN8-1], Mouse, Monoclonal |
|----------------------------|--|
| | Cytokeratin o Antibody, Cione. [CTKNO-1], Mouse, Monocional |
| Biozol Catalog Number | NSJ-V7042SAF-100UG |
| Supplier Catalog Number | V7042SAF-100UG |
| Alternative Catalog Number | NSJ-V7042SAF-100UG |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | FACS, IF, IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant human protein was used as the immunogen for this Cytokeratin 8 antibody. |
| Product Description | Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). Cytokeratin 8 is primarily found in the non-squamous epithelia and is present in majority |
| Clonality | Monoclonal |
| Concentration | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Clone Designation | [CYKN8-1] |
| Isotype | Mouse IgG1 |
| NCBI | 3856 |

| Buffer | 1X PBS, pH 7.4 |
|--------------------|---|
| Purity | Protein G affinity chromatography |
| Form | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Target | KRT8 |
| Antibody Type | Primary Antibody |
| Application Dilute | Flow cytometry: 1-2ug/10 6 cells, Western blot: 1-2ug/ml, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT, Prediluted IHC only format: incubate for 30 min at RT (1) |
| Application Notes | The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the antibody to be titered up or down for optimal performance.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |