

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

PAX7 Antibody, Clone: [PBPX7], Mouse, Monoclonal NSJ-V7047-20UG

Article Name	PAX7 Antibody, Clone: [PBPX7], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V7047-20UG
Supplier Catalog Number	V7047-20UG
Alternative Catalog Number	NSJ-V7047-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	A recombinant fragment (aa300-600) was used as the immunogen for the PAX7 antibody.
Product Description	The Pax gene family of nuclear transcription factors is comprised of nine members that function during embryogenesis to regulate the temporal and position-dependent differentiation of cells. Pax-7 is a protein specifically expressed in cultured satel
Clonality	Monoclonal
Concentration	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Clone Designation	[PBPX7]
Isotype	Mouse IgG1, kappa

UniProt	P23759
Buffer	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Purity	Protein G affinity chromatography
Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Target	PAX7
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT,Prediluted IHC only format: incubate for 30 min at RT (1)
Application Notes	Optimal dilution of the PAX7 antibody should be determined by the researcher.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.