

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

PAX7 Antibody, Clone: [PBPX7], Mouse, Monoclonal NSJ-V7047SAF-100UG

PAX7 Antibody, Clone: [PBPX7], Mouse, Monoclonal
NSJ-V7047SAF-100UG
V7047SAF-100UG
NSJ-V7047SAF-100UG
NSJ Bioreagents
Mouse
Antikörper
IHC-P
Human
A recombinant fragment (aa300-600) was used as the immunogen for the PAX7 antibody.
The Pax gene family of nuclear transcription factors is comprised of nine members that function during embryogenesis to regulate the temporal and position-dependent differentiation of cells. Pax-7 is a protein specifically expressed in cultured satel
Monoclonal
1 mg/ml in 1X PBS, BSA free, sodium azide free
[PBPX7]
Mouse IgG1, kappa
P23759

Buffer	1 mg/ml in 1X PBS, BSA free, sodium azide free
Purity	Protein G affinity chromatography
Form	1 mg/ml in 1X PBS, BSA free, sodium azide free
Target	PAX7
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT,Prediluted IHC only format: incubate for 30 min at RT (1)
Application Notes	Optimal dilution of the PAX7 antibody should be determined by the researcher.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.