

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Kappa Antibody, Clone: [KC3L-1], Mouse, Monoclonal NSJ-V7052IHC-7ML

Article Name	Kappa Antibody, Clone: [KC3L-1], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V7052IHC-7ML
Supplier Catalog Number	V7052IHC-7ML
Alternative Catalog Number	NSJ-V7052IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Human Kappa light chain was used as the immunogen for this Kappa antibody.
Product Description	This antibody is specific to the kappa light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. In mammals, the two light chains in an antibody are always identical, with only one type of ligh
Clonality	Monoclonal
Concentration	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide, *For IHC use only*
Clone Designation	[KC3L-1]
Isotype	Mouse IgG1, kappa

NCBI	3514
Buffer	1X PBS, pH 7.4
Purity	Protein G affinity chromatography
Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Target	Kappa Light Chain
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the Kappa antibody to be titered up or down for optimal performance.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.