

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

Moesin Antibody, Clone: [EDLP70], Mouse, Monoclonal NSJ-V7107SAF-100UG

Article Name	Moesin Antibody, Clone: [EDLP70], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V7107SAF-100UG
Supplier Catalog Number	V7107SAF-100UG
Alternative Catalog Number	NSJ-V7107SAF-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	FACS, IF, IHC-P, WB
Species Reactivity	Human
Immunogen	Recombinant full-length human protein was used as the immunogen for the Moesin antibody.
Product Description	Moesin (for membrane-organizing extension spike protein) is a member of the ERM protein family which includes ezrin and radixin. ERM proteins appear to function as cross-linkers between plasma membranes and actin-based cytoskeletons. Moesin is locali
Clonality	Monoclonal
Concentration	1 mg/ml in 1X PBS, BSA free, sodium azide free
Clone Designation	[EDLP70]
Isotype	Mouse IgG1, kappa
UniProt	P26038

Buffer	1 mg/ml in 1X PBS, BSA free, sodium azide free
Purity	Protein G affinity chromatography
Form	1 mg/ml in 1X PBS, BSA free, sodium azide free
Target	Moesin
Antibody Type	Primary Antibody
Application Dilute	Western blot: 1-2ug/ml,Flow Cytometry: 0.5-1ug/million cells in 0.1ml,Immunofluorescence: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT,Prediluted IHC only format: incubate for 30 min at RT (1)
Application Notes	Titering of the Moesin antibody may be required for optimal performance.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.