

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Cytokeratin 18 Antibody, Clone: [CTKN18-1], Mouse, Monoclonal NSJ-V7110SAF-100UG

Cytokeratin 18 Antibody, Clone: [CTKN18-1], Mouse, Monoclonal
NSJ-V7110SAF-100UG
V7110SAF-100UG
NSJ-V7110SAF-100UG
NSJ Bioreagents
Mouse
Antikörper
IF, IHC-P, WB
Human
Recombinant full-length human KRT18 protein was used as the immunogen for the Cytokeratin 18 antibody.
Keratin 18 is a type I cytokeratin. It is, together with its filament partner keratin 8, perhaps the most commonly found products of the intermediate filament gene family. They are expressed in single layer epithelial tissues of the body. Mutations i
Monoclonal
1 mg/ml in 1X PBS, BSA free, sodium azide free
[CTKN18-1]
Mouse IgG1, kappa
P05783

Buffer	1 mg/ml in 1X PBS, BSA free, sodium azide free
Purity	Protein G affinity chromatography
Form	1 mg/ml in 1X PBS, BSA free, sodium azide free
Target	Cytokeratin 18
Antibody Type	Primary Antibody
Application Dilute	Western blot: 1-2ug/ml,Immunofluorescence: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT,Prediluted IHC only format: incubate for 30 min at RT (1)
Application Notes	Titering of the Cytokeratin 18 antibody may be required for optimal performance.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.