

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Smooth Muscle Actin Antibody, Clone: [SMAT2-1], Mouse, Monoclonal NSJ-V7150SAF-100UG

| Article Name | Smooth Muscle Actin Antibody, Clone: [SMAT2-1], Mouse, Monoclonal |
|----------------------------|--|
| Biozol Catalog Number | NSJ-V7150SAF-100UG |
| Supplier Catalog Number | V7150SAF-100UG |
| Alternative Catalog Number | NSJ-V7150SAF-100UG |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | IF, IHC-P |
| Species Reactivity | Human, Rat |
| Immunogen | Recombinant full-length human ACTA2 protein was used as the immunogen for the Smooth Muscle Actin antibody. |
| Product Description | Actins are a family of proteins which are highly conserved and play a role in cell motility, structure and integrity. Alpha, beta and gamma actin isoforms have been identified, with alpha actins being a major constituent of the contractile apparatus, |
| Clonality | Monoclonal |
| Concentration | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Clone Designation | [SMAT2-1] |
| Isotype | Mouse IgG2a, kappa |
| UniProt | P62736 |

| Buffer | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
|--------------------|--|
| Purity | Protein G affinity chromatography |
| Form | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Target | Alpha Actin |
| Antibody Type | Primary Antibody |
| Application Dilute | Immunofluorescence: 1-2ug/ml,Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT,Prediluted IHC only format: incubate for 30 min at RT (1) |
| Application Notes | Optimal dilution of the Smooth Muscle Actin antibody should be determined by the researcher.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |