

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant MALT1 (MALT-Lymphoma Marker) Antibody, Clone: [MT1/3159R], Rabbit, Monoclonal NBT-10892-RBM2-P0

| | |
|----------------------------|---|
| Article Name | Recombinant MALT1 (MALT-Lymphoma Marker) Antibody, Clone: [MT1/3159R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-10892-RBM2-P0 |
| Supplier Catalog Number | 10892-RBM2-P0 |
| Alternative Catalog Number | NBT-10892-RBM2-P0-20,NBT-10892-RBM2-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Human MALT1 recombinant fragment (aa701-808) |
| Product Description | Mucosa associated lymphoid tissue lymphoma translocation gene 1 (MALT1) is found in extranodal low-grade B cell lymphomas. MALT1 encodes two Ig-like C2-type domains and fuses with an API2 gene, which is highly expressed in adult lymphoid tissue. The ... |
| Clonality | Monoclonal |
| Clone Designation | [MT1/3159R] |
| Molecular Weight | 93kDa |
| NCBI | 10892 |

| | |
|-------------------|---|
| UniProt | Q9UDY8 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) ,(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution |