

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Chromogranin A / CHGA (Neuroendocrine Marker) Antibody, IgG1, Clone: [LK2H10 + PHE5 + CGA/414], Mouse, Monoclonal NBT-1113-MSM5-P1ABX

Article Name	Chromogranin A / CHGA (Neuroendocrine Marker) Antibody, IgG1, Clone: [LK2H10 + PHE5 + CGA/414], Mouse, Monoclonal
Biozol Catalog Number	NBT-1113-MSM5-P1ABX
Supplier Catalog Number	1113-MSM5-P1ABX
Alternative Catalog Number	NBT-1113-MSM5-P1ABX-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human, Monkey, Mouse, Porcine, Rat
Immunogen	Human pheochromocytoma (LK2H10 & PHE5) & recombinant human chromogranin A protein (CGA414)
Product Description	Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, para
Clonality	Monoclonal
Clone Designation	[LK2H10 + PHE5 + CGA/414]
Molecular Weight	68-75kDa
Isotype	IgG1

NCBI	1113
UniProt	P10645
Form	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are also available WITHOUT BSA & Double at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Application Notes	Immunohistochemistry (Formalin-fixed) (0.1-0.2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f