

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cathepsin D (Tumor Marker) Antibody, IgG1, Clone: [CTSD/4497], Mouse, Monoclonal NBT-1509-MSM7-P1ABX

| | |
|----------------------------|---|
| Article Name | Cathepsin D (Tumor Marker) Antibody, IgG1, Clone: [CTSD/4497], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-1509-MSM7-P1ABX |
| Supplier Catalog Number | 1509-MSM7-P1ABX |
| Alternative Catalog Number | NBT-1509-MSM7-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Purified His-tagged CTSD protein |
| Product Description | Cathepsin D is a ubiquitously expressed lysosomal aspartyl protease involved in the normal degradation of proteins. It is synthesized as an inactive 43kDa preprocathepsin D that is cleaved and glycosylated to form a 46kDa procathepsin D and then furt... |
| Clonality | Monoclonal |
| Clone Designation | [CTSD/4497] |
| Molecular Weight | 48kDa (Procathepsin D), 43kDa (preprocathepsin D), 34kDa (Cathepsin D), 28kDa (heavy chain), 15kDa (light chain) |
| Isotype | IgG1 |

| | |
|-------------------|--|
| NCBI | 1509 |
| UniProt | P07339 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Western Blot (1-2ug/ml),,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 2 |