

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Desmin (Muscle Cell Marker) Antibody, IgG1, Clone: [DES/4526], Mouse, Monoclonal NBT-1674-MSM6-P1ABX

| | |
|----------------------------|---|
| Article Name | Desmin (Muscle Cell Marker) Antibody, IgG1, Clone: [DES/4526], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-1674-MSM6-P1ABX |
| Supplier Catalog Number | 1674-MSM6-P1ABX |
| Alternative Catalog Number | NBT-1674-MSM6-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human desmin protein |
| Product Description | Cytoskeletal intermediate filaments (IFs) constitute a diverse group of proteins that are expressed in a highly tissue-specific manner. IFs are constructed from two-chain α -helical coiled-coil molecules arranged on an imperfect helical lattice, and h... |
| Clonality | Monoclonal |
| Clone Designation | [DES/4526] |
| Molecular Weight | 52kDa |
| Isotype | IgG1 |

| | |
|-------------------|---|
| NCBI | 1674 |
| UniProt | P17661 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |