

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Heart Fatty Acid Binding Protein (H-FABP) / FABP3 Antibody, Clone: [FABP3/8343R], Rabbit, Monoclonal NBT-2170-RBM6-P0

| | |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name | Recombinant Heart Fatty Acid Binding Protein (H-FABP) / FABP3 Antibody, Clone: [FABP3/8343R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-2170-RBM6-P0 |
| Supplier Catalog Number | 2170-RBM6-P0 |
| Alternative Catalog Number | NBT-2170-RBM6-P0-20,NBT-2170-RBM6-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Human recombinant FABP3 protein fragment (around aa1-127) (exact sequence is proprietary) |
| Product Description | The intracellular fatty acid-binding proteins (FABPs) belongs to a multigene family. FABPs are divided into at least three distinct types, namely the hepatic-, intestinal- and cardiac-type. They form 14-15 kDa proteins and are thought to participate ... |
| Clonality | Monoclonal |
| Clone Designation | [FABP3/8343R] |
| Molecular Weight | 15kDa |
| NCBI | 2170 |

| | |
|-------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt | P05413 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |