

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant CD50 / ICAM3 Antibody, IgG1, Clone: [rICAM3/1019], Mouse, Monoclonal NBT-3385-MSM7-P0

| Article Name | Recombinant CD50 / ICAM3 Antibody, IgG1, Clone: [rlCAM3/1019], Mouse, Monoclonal |
|----------------------------|--|
| Biozol Catalog Number | NBT-3385-MSM7-P0 |
| Supplier Catalog Number | 3385-MSM7-P0 |
| Alternative Catalog Number | NBT-3385-MSM7-P0-20,NBT-3385-MSM7-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human ICAM3 protein |
| Product Description | Recognizes the D1 domain of an N-glycosylated glycoprotein of 120kDa with intra-chain disulfide bonds, identified as CD50 or ICAM-3 (Leucocyte Workshop VI). CD50 is the major ligand for LFA-1 (CD11a/CD18) and may have signaling role to increase adhes |
| Clonality | Monoclonal |
| Clone Designation | [rICAM3/1019] |
| Molecular Weight | 110-160kDa |
| Isotype | IgG1 |

| NCBI | 3385 |
|-------------------|--|
| UniProt | P32942 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a side. Also available WITHOUT BSA & Double at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Blocks binding of CD11a (LFA-1) to CD50 (ICAM-3) (Order Ab without Azide),Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 4 |