

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Recombinant Cytokeratin 17 (CK17) (Basal Epithelial Marker) Antibody, Clone: [KRT17/8320R], Rabbit, Monoclonal NBT-3872-RBM6-P1ABX

|                            |   |
|----------------------------|---|
| Article Name               | Recombinant Cytokeratin 17 (CK17) (Basal Epithelial Marker) Antibody, Clone: [KRT17/8320R], Rabbit, Monoclonal  |
| Biozol Catalog Number      | NBT-3872-RBM6-P1ABX   |
| Supplier Catalog Number    | 3872-RBM6-P1ABX   |
| Alternative Catalog Number | NBT-3872-RBM6-P1ABX-100   |
| Manufacturer               | NeoBiotechnologies  |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | IHC   |
| Species Reactivity         | Human   |
| Immunogen                  | Recombinant fragment of human KRT17 protein (exact sequence is proprietary)   |
| Product Description        | Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excellent tool to distinguish myoepithelial cells from luminal epithelium of various glandsvc OOsuch... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [KRT17/8320R]   |
| Molecular Weight           | 46kDa   |
| NCBI                       | <a href="#">3872</a>  |

|                   |  |
|-------------------|--|
| UniProt           | <a href="#">Q04695</a>   |
| Form              | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type     | Recombinant Monoclonal Antibody  |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes),Optimal dilutio |