

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Cytokeratin 19 (KRT19) (Pancreatic Stem Cell Marker) Antibody, Clone: [KRT19/1959R], Rabbit, Monoclonal NBT-3880-RBM9-P0

| | |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name | Recombinant Cytokeratin 19 (KRT19) (Pancreatic Stem Cell Marker) Antibody, Clone: [KRT19/1959R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-3880-RBM9-P0 |
| Supplier Catalog Number | 3880-RBM9-P0 |
| Alternative Catalog Number | NBT-3880-RBM9-P0-20,NBT-3880-RBM9-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant human full-length KRT19 protein |
| Product Description | This Ab reacts with the rod domain of human cytokeratin 19 (CK19), a polypeptide of 40kDa. CK19 is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, ... |
| Clonality | Monoclonal |
| Clone Designation | [KRT19/1959R] |
| Molecular Weight | 40kDa |
| NCBI | 3880 |

| | |
|-------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt | P08727 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT), (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minute |