

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

PAX3 Antibody, IgG1, Clone: [PAX3/8426], Mouse, Monoclonal NBT-5077-MSM6-P0

| | |
|----------------------------|---|
| Article Name | PAX3 Antibody, IgG1, Clone: [PAX3/8426], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-5077-MSM6-P0 |
| Supplier Catalog Number | 5077-MSM6-P0 |
| Alternative Catalog Number | NBT-5077-MSM6-P0-20,NBT-5077-MSM6-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment (around aa150-350) of human PAX3 protein (exact sequence is proprietary) |
| Product Description | Pax genes contain paired domains that share strong homology to genes in Drosophila which are involved in programming early development. The product of the PAX3 gene is a DNA-binding protein expressed during early neurogenesis. Pax-3 is a protein cont... |
| Clonality | Monoclonal |
| Clone Designation | [PAX3/8426] |
| Molecular Weight | 56kDa |
| Isotype | IgG1 |
| NCBI | 5077 |

| | |
|-------------------|--|
| UniProt | P23760 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |