

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant FOXP3 (Forkhead Box Protein P3) / Scurfin Antibody, Clone: [FOXP3/8015R], Rabbit, Monoclonal NBT-50943-RBM6-P1ABX

| | |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name | Recombinant FOXP3 (Forkhead Box Protein P3) / Scurfin Antibody, Clone: [FOXP3/8015R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-50943-RBM6-P1ABX |
| Supplier Catalog Number | 50943-RBM6-P1ABX |
| Alternative Catalog Number | NBT-50943-RBM6-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment (around aa300-431) of human FOXP3 protein (exact sequence is proprietary) |
| Product Description | Recognizes a protein of 47-55kDa, which is identified as FOXP3. Its precise epitope is not known, but it has been mapped to the N-terminal portion of the protein. The FOX family of transcription factors is a large group of proteins that share a commo... |
| Clonality | Monoclonal |
| Clone Designation | [FOXP3/8015R] |
| Molecular Weight | 47-55kDa |
| NCBI | 50943 |

| | |
|-------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt | Q9BZS1 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |