

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Monocyte Chemotactic Protein 2 (MCP2) / CCL8 Antibody, IgG1, Clone: [CCL8/3686], Mouse, Monoclonal NBT-6355-MSM6-P0

| | |
|----------------------------|---|
| Article Name | Monocyte Chemotactic Protein 2 (MCP2) / CCL8 Antibody, IgG1, Clone: [CCL8/3686], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-6355-MSM6-P0 |
| Supplier Catalog Number | 6355-MSM6-P0 |
| Alternative Catalog Number | NBT-6355-MSM6-P0-20,NBT-6355-MSM6-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Human recombinant MCP2 / CCL8 protein fragment (around aa1-99) (exact sequence is proprietary) |
| Product Description | This antimicrobial gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfam... |
| Clonality | Monoclonal |
| Clone Designation | [CCL8/3686] |
| Molecular Weight | 11kDa |
| Isotype | IgG1 |

| | |
|-------------------|--|
| NCBI | 6355 |
| UniProt | P80075 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |