

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant CD162 (Selectin P Ligand) Antibody, Clone: [PSGL1/8192R], Rabbit, Monoclonal NBT-6404-RBM5-P1ABX

| | |
|----------------------------|---|
| Article Name | Recombinant CD162 (Selectin P Ligand) Antibody, Clone: [PSGL1/8192R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-6404-RBM5-P1ABX |
| Supplier Catalog Number | 6404-RBM5-P1ABX |
| Alternative Catalog Number | NBT-6404-RBM5-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human SELPLG (CD162) protein |
| Product Description | CD162 glycoprotein functions as a high affinity counter-receptor for the cell adhesion molecules P-, E- and L- selectin expressed on myeloid cells and stimulated T lymphocytes. As such, this protein plays a critical role in leukocyte trafficking duri... |
| Clonality | Monoclonal |
| Clone Designation | [PSGL1/8192R] |
| Molecular Weight | 43-120kDa |
| NCBI | 6404 |

| | |
|-------------------|--|
| UniProt | Q14242 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |