

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant TNF-alpha (Tumor Necrosis Factor alpha) Antibody, Clone: [TNF/1500R], Rabbit, Monoclonal NBT-7124-RBM13-P1ABX

| | |
|----------------------------|---|
| Article Name | Recombinant TNF-alpha (Tumor Necrosis Factor alpha) Antibody, Clone: [TNF/1500R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-7124-RBM13-P1ABX |
| Supplier Catalog Number | 7124-RBM13-P1ABX |
| Alternative Catalog Number | NBT-7124-RBM13-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human, Rat |
| Immunogen | Recombinant full-length human TNF-alpha protein |
| Product Description | This MAb recognizes human 17-26kDa protein, which is identified as cytokine TNF-alpha (Tumor Necrosis Factor-alpha). TNF-alpha can be expressed as a 17kDa free molecule, or as a 26kDa membrane protein. TNF-alpha is a protein secreted by lipopolysacch... |
| Clonality | Monoclonal |
| Clone Designation | [TNF/1500R] |
| Molecular Weight | 17kDa |
| NCBI | 7124 |

| | |
|-------------------|---|
| UniProt | P01375 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (4-8ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution for a |