

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Vimentin (Mesenchymal Cell Marker) Antibody, IgM, Clone: [LN-6], Mouse, Monoclonal NBT-7431-MSM2-P1BX

| | |
|----------------------------|---|
| Article Name | Vimentin (Mesenchymal Cell Marker) Antibody, IgM, Clone: [LN-6], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-7431-MSM2-P1BX |
| Supplier Catalog Number | 7431-MSM2-P1BX |
| Alternative Catalog Number | NBT-7431-MSM2-P1BX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Bovine, Feline, Human, Mouse, Porcine, Rabbit, Rat, Sheep |
| Immunogen | Human thymic nuclear extract |
| Product Description | This MAb reacts with a 58kDa protein identified as vimentin. It reacts with a non-hematopoietic epitope of vimentin and shows no cross-reaction with other closely related intermediate filament proteins (IFPs) such as desmin, keratin, neurofilament, a... |
| Clonality | Monoclonal |
| Clone Designation | [LN-6] |
| Molecular Weight | 57-60kDa |
| Isotype | IgM |

| | |
|-------------------|---|
| NCBI | 7431 |
| UniProt | P08670 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (0.1-0.2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f |