

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) Antibody, IgG1, Clone: [3E2D10 + VWF635], Mouse, Monoclonal NBT-7450-MSM3-P1ABX**

Article Name	von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) Antibody, IgG1, Clone: [3E2D10 + VWF635], Mouse, Monoclonal
Biozol Catalog Number	NBT-7450-MSM3-P1ABX
Supplier Catalog Number	7450-MSM3-P1ABX
Alternative Catalog Number	NBT-7450-MSM3-P1ABX-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC, WB
Species Reactivity	Human
Immunogen	Recombinant human vWF fragment spanning around aa 845-949 (exact sequence is proprietary)
Product Description	von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation...
Clonality	Monoclonal
Clone Designation	[3E2D10 + VWF635]
Molecular Weight	250kDa
Isotype	IgG1

NCBI	<a href="#">7450</a>
UniProt	<a href="#">P04275</a>
Form	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Application Notes	Immunoprecipitation (1-2 g/500ug protein lysate), Flow Cytometry (1-2ug/million cells), ,Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues re