

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) Antibody, IgG2b, Clone: [VWF/1767], Mouse, Monoclonal NBT-7450-MSM7-P1ABX**

Article Name	von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) Antibody, IgG2b, Clone: [VWF/1767], Mouse, Monoclonal
Biozol Catalog Number	NBT-7450-MSM7-P1ABX
Supplier Catalog Number	7450-MSM7-P1ABX
Alternative Catalog Number	NBT-7450-MSM7-P1ABX-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	ELISA, IHC
Species Reactivity	Human
Immunogen	Recombinant fragment (around aa1815-1939) of human vWF protein (exact sequence is proprietary)
Product Description	von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational...
Clonality	Monoclonal
Clone Designation	[VWF/1767]
Molecular Weight	250kDa
Isotype	IgG2b

NCBI	<a href="#">7450</a>
UniProt	<a href="#">P04275</a>
Form	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Application Notes	ELISA (Use Ab at 2-4ug/ml for coating) (Order Ab without BSA), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95