

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant CD3e (T-Cell Marker) Antibody, IgG1, Clone: [rC3e/1308], Mouse, Monoclonal NBT-916-MSM19-P1ABX

| | |
|----------------------------|---|
| Article Name | Recombinant CD3e (T-Cell Marker) Antibody, IgG1, Clone: [rC3e/1308], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-916-MSM19-P1ABX |
| Supplier Catalog Number | 916-MSM19-P1ABX |
| Alternative Catalog Number | NBT-916-MSM19-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human CD3E protein |
| Product Description | Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the... |
| Clonality | Monoclonal |
| Clone Designation | [rC3e/1308] |
| Molecular Weight | 20kDa |
| Isotype | IgG1 |

| | |
|-------------------|---|
| NCBI | 916 |
| UniProt | P07766 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f |