

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) Antibody, Clone: [Ki-1/1747R], Rabbit, Monoclonal NBT-943-RBM7-P1ABX

| | |
|----------------------------|---|
| Article Name | Recombinant CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) Antibody, Clone: [Ki-1/1747R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-943-RBM7-P1ABX |
| Supplier Catalog Number | 943-RBM7-P1ABX |
| Alternative Catalog Number | NBT-943-RBM7-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant human full-length TNFRSF8 protein |
| Product Description | Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease,... |
| Clonality | Monoclonal |
| Clone Designation | [Ki-1/1747R] |
| Molecular Weight | 105-120kDa |
| NCBI | 943 |

| | |
|-------------------|---|
| UniProt | P28908 |
| Form | 200ug/ml of Ab Purified by Protein A Column. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 8.0, for 45 min at 95°C followed by cooling at RT for 20 minutes)Optimal dilution |