

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant CD44v4 (Marker of Tumor Metastasis) Antibody, IgG2a, Clone: [rCD44v4/1219], Mouse, Monoclonal NBT-960-MSM12-P0

| | |
|----------------------------|---|
| Article Name | Recombinant CD44v4 (Marker of Tumor Metastasis) Antibody, IgG2a, Clone: [rCD44v4/1219], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-960-MSM12-P0 |
| Supplier Catalog Number | 960-MSM12-P0 |
| Alternative Catalog Number | NBT-960-MSM12-P0-20,NBT-960-MSM12-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment corresponding to the v4 domain of human CD44 |
| Product Description | This antibody recognizes an epitope encoded by exon v4 on the variant portion of human CD44. The CD44 molecule belongs to a family of cellular adhesion molecules found on a wide range of normal and malignant cells in epithelial, mesothelial and hemop... |
| Clonality | Monoclonal |
| Clone Designation | [rCD44v4/1219] |
| Molecular Weight | 80-95kDa |
| Isotype | IgG2a |

| | |
|-------------------|---|
| NCBI | 960 |
| UniProt | P16070 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f |