

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD63 (Late Endosomes Marker) Antibody, IgG2a, Clone: [LAMP3/2881], Mouse, Monoclonal NBT-967-MSM11-P0

| | |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name | CD63 (Late Endosomes Marker) Antibody, IgG2a, Clone: [LAMP3/2881], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-967-MSM11-P0 |
| Supplier Catalog Number | 967-MSM11-P0 |
| Alternative Catalog Number | NBT-967-MSM11-P0-20,NBT-967-MSM11-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment (around aa 100-197) of human CD63 protein |
| Product Description | This MAAb recognizes protein of 26kDa-60kDa, which is identified as CD63. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with spec... |
| Clonality | Monoclonal |
| Clone Designation | [LAMP3/2881] |
| Molecular Weight | 26kDa (core protein), 30-60kDa (glycosylated) |
| Isotype | IgG2a |

| | |
|-------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| NCBI | 967 |
| UniProt | P08962 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minu |