

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD79b (B-Cell Marker) Antibody, IgG1, Clone: [IGB/2555], Mouse, Monoclonal NBT-974-MSM5-P0

| | |
|----------------------------|---|
| Article Name | CD79b (B-Cell Marker) Antibody, IgG1, Clone: [IGB/2555], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-974-MSM5-P0 |
| Supplier Catalog Number | 974-MSM5-P0 |
| Alternative Catalog Number | NBT-974-MSM5-P0-20,NBT-974-MSM5-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | ELISA, IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment of human CD79b protein (around aa 29-159) (exact sequence is proprietary) |
| Product Description | CD79 (also designated Ig chains, designated CD79B or B29. The B cell antigen receptor complex (BCR) is formed by the association of CD79 with a membrane immunoglobulin, such as IgM or IgD. The membrane immunoglobulins IgM and IgD achieve surface expr... |
| Clonality | Monoclonal |
| Clone Designation | [IGB/2555] |
| Molecular Weight | 39kDa |
| Isotype | IgG1 |

| | |
|-------------------|--|
| NCBI | 974 |
| UniProt | P40259 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | ELISA (Use Ab at 2-4ug/ml for coating) (Order Ab without BSA),,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 |