

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Bromodeoxyuridine (BrdU) (Proliferation Marker) Antibody, IgG1, Clone: [MoBu-1], Mouse, Monoclonal NBT-MSM4-884-P0

| | |
|----------------------------|---|
| Article Name | Bromodeoxyuridine (BrdU) (Proliferation Marker) Antibody, IgG1, Clone: [MoBu-1], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-MSM4-884-P0 |
| Supplier Catalog Number | MSM4-884-P0 |
| Alternative Catalog Number | NBT-MSM4-884-P0-20,NBT-MSM4-884-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC |
| Species Reactivity | All |
| Immunogen | Bromodeoxyuridine (BrdU) conjugated to hemocyanine (isolated from <i>Helix pomatia</i>) |
| Product Description | It reacts with Bromodeoxyuridine (BrdU) in single stranded DNA (produced by partial denaturation of double stranded DNA), BrdU coupled to a protein carrier, as well as free BrdU. BrdU is a thymidine analog, incorporated into cell nuclei during DNA sy... |
| Clonality | Monoclonal |
| Clone Designation | [MoBu-1] |
| Molecular Weight | Depends on the target |
| Isotype | IgG1 |

| | |
|-------------------|--|
| NCBI | Not Applicable |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) [For staining of formalin-fixed tissues, incubate sections in 4N HCl for 30 minutes at RT followed by digestion wit |