

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Cytokeratin, Acidic (Type I or LMW) (Epithelial Marker) Antibody, Clone: [KRTL/1577R], Rabbit, Monoclonal NBT-RBM21-1577-P0

Article Name	Recombinant Cytokeratin, Acidic (Type I or LMW) (Epithelial Marker)
	Antibody, Clone: [KRTL/1577R], Rabbit, Monoclonal
Biozol Catalog Number	NBT-RBM21-1577-P0
Supplier Catalog Number	RBM21-1577-P0
Alternative Catalog Number	NBT-RBM21-1577-P0-20,NBT-RBM21-1577-P0-100
Manufacturer	NeoBiotechnologies
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Canine, Human
Immunogen	Recombinant human KRT77 protein fragment
Product Description	This MAb recognizes the 56.5kDa (CK10), 50kDa (CK14), 50kDa (CK15), 48kDa (CK16), 40kDa (CK19) keratins of the acidic (Type I or LMW) subfamily. Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pl 6.0) subfamil
Clonality	Monoclonal
Clone Designation	[KRTL/1577R]
Molecular Weight	56.5kDa (CK10), 50kDa (CK14), 50kDa (CK15), 48kDa (CK16), 40kDa (CK19)
NCBI	374454

UniProt	Q7Z794
Form	200ug/ml of Ab produced in HEK293 cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & amp; 0.05% azide. Also available WITHOUT BSA & amp; azide at 1.0mg/ml.
Antibody Type	Recombinant Monoclonal Antibody
Application Notes	Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution for a