

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Cytokeratin, pan (Epithelial Marker) Antibody, Clone: [Cocktail PCK/4933R], Rabbit, Monoclonal NBT-RBM43-4933-P1ABX

| | |
|----------------------------|---|
| Article Name | Recombinant Cytokeratin, pan (Epithelial Marker) Antibody, Clone: [Cocktail PCK/4933R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-RBM43-4933-P1ABX |
| Supplier Catalog Number | RBM43-4933-P1ABX |
| Alternative Catalog Number | NBT-RBM43-4933-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Crude cytokeratin extract prepared from RT-4 and MCF-7 cells |
| Product Description | PCK/4933R demonstrates a broad spectrum of cytokeratin reactivity. In normal tissues, PCK/4933R is reactive with most epithelial types, including bile ducts and hepatocytes in liver, bladder epithelium, breast ducts, bronchial epithelium, endometrium... |
| Clonality | Monoclonal |
| Clone Designation | [Cocktail PCK/4933R] |
| Molecular Weight | 40-67kDa (Multiple) |
| NCBI | Multiple |

| | |
|-------------------|---|
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |