



Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## M22, CAS [[864420-54-2]] FBM-10-5733

| Article Name               | M22, CAS [[864420-54-2]]          |
|----------------------------|-----------------------------------|
| Biozol Catalog Number      | FBM-10-5733                       |
| Supplier Catalog Number    | 10-5733                           |
| Alternative Catalog Number | FBM-10-5733-5MG,FBM-10-5733-25MG  |
| Manufacturer               | Focus Biomolecules                |
| Category                   | Biochemikalien                    |
| Product Description        | NEDD8 activating enzyme inhibitor |
| Molecular Weight           | 363.33                            |
| Purity                     | >98% by TLC NMR (Conforms)        |
| Form                       | Off-white solid                   |
| CAS Number                 | [864420-54-2]                     |
| Formula                    | C20H24Cl2N2                       |

**Application Notes** 

Protein neddylation is a post-translational modification which transfers the ubiquitin-like protein NEDD8 to a lysine residue of the target substrate through a three-step enzymatic cascade and has been considered as a viable target for the development of new therapeutics for cancer.1 M22 (CAS 864420-54-2) is an inhibitor of NEDD8 activating enzyme (NAE) which was discovered by virtual screening of a small molecule library (50,000 compounds) against the active site of NAE.2 M22 is selective and reversible and inhibits NEDDylation of ubiquitin-conjugating enzyme 12 (UBC12) in a cell-free assay employing recombinant human NAE. It selectively inhibits NEDDylation over SUMOylation and ubiquitination in A549 cells and inhibits their growth (GI50 = 5.5  $\mu$ M). It has been shown to inhibit tumor growth in a mouse xenograft model (at 60 mg/kg).2