

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Biotinylated human FcRn (FCGRT-B2M) heterodimer protein GMP-HUFCRN-B

Article Name	Biotinylated human FcRn (FCGRT-B2M) heterodimer protein
Biozol Catalog Number	GMP-HUFCRN-B
Supplier Catalog Number	HUFCRN-B
Alternative Catalog Number	GMP-HUFCRN-B-250, GMP-HUFCRN-B-5X250
Manufacturer	Gamma Proteins UK
Category	Proteine/Peptide
Application	BLI, ELISA, SBR
Species Reactivity	Human
Conjugation	Biotin
Product Description	IgG Fc fragment receptor transporter, also known as the neonatal Fc receptor or more commonly FcRn, consist of two subunits (FCGRT and B2M) and forms an MHC class I-like heterodimer. FcRn binds to the Fc domain of monomeric IgG and mediates the pH de
Molecular Weight	46,6 kDa
UniProt	P55899
Buffer	PBS, pH7.4
Source	The sequence of the extracellular domain of human FCGRT (Ala 24-Ser 297) was fused with a C-terminal tag consisting of the AVI tag, TEV protease recognition sequence and a 10-His tag. This was cotransfected with the sequence of human B2M (Ile 21-Met 119)
Expression System	HEK293

Purity	>95 % as determined by SEC-HPLC
Form	Lyophilized
Sequence	A24-S297 & amp; I21-M119